



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

GENERAL NOTES.

Nesting of Man-o'-war-bird (*Fregata aquila*) in Cuba.— Puerto Escondido is a small and from the open sea invisible port on the south coast of Cuba a few miles to the east of Guantanamo Bay. It was used in the old days as a pirate retreat, being admirably fitted for the purpose by its narrow entrance and inconspicuousness, and is so mentioned in 'Tom Cringles Log' by Mr. Scott. The name Puerto Escondido means the Hidden Port. It will now form part of the U. S. Naval Station at Guantanamo Bay and will therefore be American territory. In this port is a small Key — Cayo Guincho — about 100 feet in diameter composed mostly of mud upon which grows, in the center, a large mangroove tree surrounded by smaller ones in the water. Upon approaching the key on Dec. 23, 1910, large numbers of Man-o'-war-birds were seen flying about it and upon a closer inspection it was noticed that the large tree had a number of nests upon it. The birds, frightened by the launch, left the nests so hurriedly that some eggs were knocked out, while below, among the roots of the smaller mangroves in the water, two species of "snapper" (*Neomænis cyanopterus* Linn. and *Neomænis griseus* Linn.) were greedily feasting on the fallen eggs.— CHARLES T. RAMSDEN, *Guantanamo, Cuba*.

American Merganser in South Carolina.— On January 2, 1911, I secured the first authentic record for the American Merganser (*Mergus americanus*) in this State. Mr. Arthur T. Wayne, in his 'Birds of South Carolina,' refers this species to a hypothetical list, and declares that to the best of his knowledge it has never been taken in the State. The specimen here recorded was shot on Back River near its junction with the Cooper, after having partially swallowed a mullet ten inches in length. The weight of the mullet in its throat destroyed its balance and made continuous flight impossible. The bird rose half a dozen times but with each attempt at escape could gain only a few yards.

Several years ago a Merganser was taken on the upper Cooper River which I afterwards decided was the *Mergus americanus*, but as this specimen was not preserved I have been on the lookout during the past season for another to prove the occurrence of the species in South Carolina.

The specimen taken on Back River is a fine adult male in full plumage with the breast and under-parts a delicate peach-blossom color. The skin is now in the collection of the Charleston Museum (Spec. No. 7258).— CASPAR CHISOLM, *Charleston, S. C.*

Additional Records of the European Widgeon (*Mareca penelope*).— The last captures for the interior I recorded in 'The Auk' (Vol. XXII, 1905, p. 206). As a matter of convenient reference I now add to the list three specimens.